

## Sources, extraction and biological activities of cinnamaldehyde

Anna Ashehe Ibi<sup>1,\*</sup>, Christal Kabele Kyuka<sup>1,2</sup>

<sup>1</sup>Biotechnology Advanced Research Centre, Sheda Science and Technology Complex, P.M.B 186, Garki, Abuja, Nigeria

<sup>2</sup>Department of Crop and Horticultural Science, Pan African University for Life and Earth Sciences Institute (Including Health and Agriculture), University of Ibadan, Nigeria.

### Abstract

Cinnamaldehyde is the prevalent bioactive part of cinnamon essential oil which is liable for its regular scent and can be gotten from the bark, leaves, and twigs of various *Cinnamomum* species. Cinnamaldehyde is known to be generally considered non-toxic due to its high tolerance in animals and humans. Various extraction methods have been used for extracting cinnamaldehyde and different phytochemicals from plants. The methods generally adopted for cinnamaldehyde extraction are hydro distillation, supercritical carbon dioxide extraction, ultrasound-assisted extraction, microwave-assisted hydro distillation, and water steam extraction. Cinnamaldehyde has been documented to have different useful properties against disease conditions as a result of oxidative stress, inflammation, loss of neurons, hyperglycemia, and malignant growth. Likewise, cinnamaldehyde has been recorded to possess strong antimicrobial activity against a wide range of pathogenic and food waste microorganisms. The mechanism through which cinnamaldehyde exerts these effects have been associated with the prevention of reactive oxygen species and reactive nitrogen species generation, free radical scavenging activity, inhibiting inflammatory cytokines, and disruption of the cell membrane of microorganisms. This article gives a thorough report of the sources, extraction techniques, validated therapeutic potentials and mechanisms of cinnamaldehyde. It likewise features other applications of cinnamaldehyde in agriculture, food, and other industries.

**Keywords:** Antimicrobial, Anticancer, Antidiabetes, Cinnamon, Cinnamaldehyde.

Please cite this article as: Anna Ashehe Ibi, Christal Kabele Kyuka. Sources, extraction and biological activities of cinnamaldehyde. Trends in Pharmaceutical Sciences. 2022;8(4):263-282. doi: 10.30476/TIPS.2022.96263.1160

### Introduction

The use of medicinal plants by the traditional system for treatment and management of diseases continues to be the main stay in low-income countries due to their natural source, safety, affordability, and availability (1, 2). Unpleasant side effects and the cost of available drugs are underlying reasons, among others, which have triggered a renewed interest in natural sources for the development of therapeutic agents. Extracts, essential oils and other plant derivatives from various parts of plants are different forms in which these plants are

used for their therapeutic potentials. These pharmacological effects of medicinal plants, such as antimalarial, anticancer, antifungal, antiviral and antidiabetes, among others, are because of the inherent secondary metabolites known as phytochemicals (3). The most common phytochemicals such as flavonoids, phenols, sterols, tocopherols, and organic acids possess appreciable health benefits and other applications in the development of pharmacotherapeutic and nutraceutical foods. This review article lays emphasis on the therapeutic potential of cinnamaldehyde in the management of commonly occurring diseases and other applications.

Cinnamaldehyde also known as cinnamic aldehyde is an  $\alpha$ ,  $\beta$ -unsaturated aldehyde derived

*Corresponding Author:* Anna Ashehe Ibi, Biotechnology Advanced Research Centre, Sheda Science and Technology Complex, P.M.B 186, Garki, Abuja, Nigeria.  
Email: asheyibi@gmail.com

**Table 1.** General physicochemical properties of cinnamalydehyde.

Properties	Description
IUPAC name	(E)-3-phenylprop-2-enal
Chemical formula	C <sub>9</sub> H <sub>8</sub> O
Molecular mass	132.16 g/mol
Solubility in water	Slightly at about 1.1g/L at 20oC
Colour	Yellow to greenish yellow
Density	1.05 g/mL at 25oC
Boiling point	248oC (478oF)
Melting point	-7.5oC (18oF)
Bisulfide solubility	100%
Purity	>98%

from cinnamon trees and other species of the genus *Cinnamomum*, it is responsible for their characteristic natural flavour and fragrance. Cinnamaldehyde is a clear yellow to greenish yellow liquid compound with a pungent spicy note and accounts for 85% of the constituents of essential oil of cinnamon bark. Cinnamaldehyde is well-tolerated in both humans and animals and generally considered as a safe bioactive. Cinnamaldehyde was first isolated from cinnamon essential oil in 1834 and was first synthesized in the laboratory in 1856 by Chiozza (4). Different methods for laboratory synthesis of cinnamaldehyde are currently available but the use of steam distillation of the oil from cinnamon bark is considered most economical (5). It can be synthesized from cinnamyl alcohol, aldol condensation of benzaldehyde and acetaldehyde etc. the general physicochemical properties of cinnamaldehyde is discussed in table 1.

Studies have shown that cinnamaldehyde possess the ability to eliminate natural or chemical toxins like ochratoxins A, as well as improve general human wellbeing (6, 7). The bioactivities of cinnamaldehyde have been duly documented in literatures (8-11) and have also been reported to have other applications in agriculture as a preservative (12). Due to the bioactivities and other applications, it is of significant worth to rationally put together research findings related to the therapeutic potentials and agricultural importance of cinnamaldehyde in other to expound its importance to human health and food security. This review will also discuss the mechanisms involved in the functionality of cinnamaldehyde preventing numerous

lifestyle related health conditions.

### Sources of Cinnamaldehyde


Cinnamaldehyde (C<sub>6</sub>H<sub>5</sub>CH=CHCHO), the major organic compound of cinnamon essential oil (13). Cinnamaldehyde, which is an important flavour component of cinnamon, is obtained from the *Cinnamomum* genus. They include *Cinnamomum zeylanicum* L (Cinnamon) (13), *Cinnamomum tamala*, *Cinnamomum pauciflorum* (14), *Cinnamomum ceylanicum* (Ceylon) which is the most appraised source, *C. cassia* (cassia) and *C. burmannii* (cassia vera) which are frequently used as cheaper alternatives (15). Cinnamaldehyde can be extracted from different parts of the plants including the bark, leaves and twigs (16, 17).

### Extraction and Purification of cinnamadehyde

The extraction and isolation cinnamaldehyde have been done in various ways since time past. This review is going to highlight some ways cinnamaldehyde is being isolated and purified as reported by several authors. Figure 1 gives a glance of the different extraction methods used for cinnamaldehyde extraction.

### High Performance Counter-Current Chromatography

In this process of isolation of cinnamaldehyde, the cinnamaldehyde will be enriched by Counter-Current Chromatography (CCC) with reversed phase mode using n-hexane-ethyl acetate-methanol-water (1:1:1:1, v/v/v/v) as the solvent system. Counter-current chromatography (ccc) is



**Figure 1.** Methods for cinnamaldehyde extraction.

one of the liquid-liquid separation methods with many technical merits. Examples include ability to take less time and lower volume of solution selective and it gains higher recoveries. Fan *et al.*, (18) reportedly used the high-performance counter-current chromatography method to extract cinnamaldehyde from cinnamon bark.

### Pressure Drop Fractional distillation

Isolation of cinnamaldehyde can also be done using the pressure drop fractionation as reported by Moreta *et al.*, (19). In the separation of cinnamaldehyde from cinnamon oil using fractional distillation, the use of pressure reduction is intended so that cinnamaldehyde can evaporate faster than regular boiling points. The pressure used in this process is usually around 29.8-30.1 mmHg and temperature 30-300 °C.

### Bisulfite addition method

Bisulfite addition method is another method used to extract and isolated cinnamaldehyde. It is carried out by reacting cinnamon oil to the saturated sodium bisulfite solution with a ratio of 1:1. The precipitate formed was further neutralized with 10% HCl solution. Furthermore, cinnamaldehyde will be extracted from the solution using petroleum ether twice and washed by sterile water. The petroleum ether extract will be applied in the evaporator (20).

### Hydro Distillation (HD)

Hydro distillation is the most widely used method for cinnamaldehyde extraction, with advantages of simple process, lowcost, and absence of solvent residue, but it gives low yield at around 1-2 % (21, 22). Hydro distillation method has long been exploited in the extraction of essential oils and bioactive components of plant materials. Hydro distillation method of extraction can be carried out in three different ways: water distillation, water and steam distillation, and direct steam distillation (23). Three main physicochemical processes involved in hydro distillation are hydrodiffusion, hydrolysis, and decomposition by heat. Hot water and steam is generally considered as the basic media to break through plant matrix, thereby releasing bioactive compounds. The indirect cooling condenses the mixture of steam and bioactives and make separation of oil and bioactive components from the water to occur. Sodium sulphate is usually used for the dry of essential oils and oil-based bioactive compounds. Since hydro distillation is often conducted at temperature above boiling point of water, some volatile components, natural pigments, and heat-labile bioactive compounds may be lost (24).

Steps in extraction of cinnamaldehyde using this method is thus; weigh 10 gram of cinnamon bark powder and mix with 100 ml distilled water. The mixture is allowed for 2 days as soaking period. Extraction of essential oil from cin-

namon bark powder then becomes easy after a soaking period. The mixture is then transferred into 2 litre distillation flask and heated on 50 °C. Distillate is condensed and collected into the beaker and then transferred into separating funnel. The mixture is then allowed to separate into oil and water phase. Oil will be settled on the water phase. Separation continues until there is no oil left in the water phase. Then store it in closed vessel at low temperature. Researchers have reportedly used this method to extract cinnamaldehyde (22, 25-29).

### Water Steam Distillation

Water steam distillation is one of the types of hydro distillation methods used in extracting and isolating cinnamaldehyde. This was reported by Alsalim *et al.*, (30) where they obtained cinnamaldehyde after 6 hours using steam distillation process following the process of Senhaji *et al* (31). In this method the cinnamon oil is usually transferred to a separation funnel, then about 5-10 ml of dichloromethane will be added to the separation funnel, shaken, and allowed to separate for 10 minutes, then dichloromethane will be drained off. This step is repeated once more. The above layer which contains the cinnamaldehyde will be dried by adding sodium sulfate until it is free flowing. The solution will then be transferred to a round bottom flask and the solid sodium sulphate was rinsed with a little more dichloromethane, then evaporated by rotary evaporator, cinnamaldehyde can then collected in dark container.

### Subcritical Water Extraction (SWE)

Subcritical water extraction (SWE) is arising as a famous option for extraction of bioactive components. Water is used as extraction solvent at temperatures between 100 °C and 374 °C with a pressure (1-22.1 MPa) high enough to maintain the liquid state (32). Under subcritical conditions, the dielectric constant of water (directly related to its polarity) is reduced with increasing extraction temperature (for example, 80.4 at 20 °C to 27 at 250 °C), which consequently decreases the water polarity. As a result, water behaves like an organic solvent, thus allowing the extraction of different classes of compounds and selective extraction

of target compounds (33). Subcritical water has been used to extract essential oil, polyphenols, and polysaccharide, *etc.* (34-41). Jayawardena and Smith (42) first extracted essential oils from cinnamon bark and leaves by SWE and confirmed that SWE is a practical method for cinnamon oil extraction. Subsequently, Eikani and colleagues (43) expanded the achievements of Jayawardena and Smith in terms of operating conditions and optimization procedure. They successfully optimized the SWE conditions of cinnamon bark oil and found that SWE is an effective method for the selective extraction of oxygenated components, such as E-cinnamaldehyde. Earlier studies (44) on the SWE of cinnamon oil also revealed that the extraction time of SWE is a sixth that of SD, and the yields of cinnamon oil and cinnamaldehyde are increased by 15.8% and 28.4%, respectively. In the mean time, it was found that the drawbacks of low mass transfer efficiency and uneven extraction became obvious alongside increase in loaded quantity and height of raw materials accumulation, particularly the material in the center of the extraction tank.

### Ultrasound Subcritical Water Extraction (USWE)

To further improve the extraction yield, ultrasound, as an extraction strengthening technology, has been introduced into supercritical fluid to extract adlay oil, passion fruit seed oil and other bioactive components (45-47). These independent studies have proved that ultrasound can enhance supercritical fluid extraction. In recent years, the application of ultrasound in the SWE system is also developing gradually and has been found to effectively increase the extraction rate of SWE (48).

The apparatus for ultrasonic subcritical water extraction consists of SWE system, ultrasonic strengthening device and operating system, which has been reported in literature (49). The ultrasonic strengthening device mainly includes ultrasonic generator and transducer. The ultrasonic output power is between 0 and 250 W. The ultrasonic frequency can be adjusted between 15 kHz and 38 kHz. This operation is like SWE, except that USWE requires adjustment of the ultrasonic strengthening device. The powdered cinnamon

bark will be extracted with distilled water at designed temperature, extraction time, pressure, and ultrasound conditions by the above USWE apparatus. After the extraction, the extract will be rapidly cooled down through cooling system to prevent degradation. Guo (50) also reported using this extraction method to extract and isolate cinnamaldehyde.

### Soxhlet distillation

This method was initially designed by German chemist Franz Ritter Von Soxhlet in 1879 for lipid extraction from a solid material. These days, it is often utilized as a reference technique to evaluate extraction yield of lipid from plant materials. The principle of Soxhlet extraction is using a solvent vapour to dissolve oil-base compounds inside the plant sample housed in a thimble. The vapour is then condensed to separate the compounds of interest from the solvent. Although Soxhlet extraction is simple, inexpensive, and easy to operate, it requires a large amount of solvent, long extraction time, and leads to degradation of heat-labile bioactive compounds (51, 52). Masghati and Ghoreishi, (53) reported using soxhlet distillation method for cinnamaldehyde extraction.

### Ultrasonic Assisted Extraction

The principle of ultrasonic assisted extraction is to use bubble cavitation created by ultrasound waves (20 kHz to 100 MHz) to rupture plant cell walls and intact plant materials, and thus, enhance the extraction efficiency. Cavitation is a phenomenon brought about by the passage of ultrasound wave through fluid medium creating alternative pressure and expansion pattern of bubbles. As the bubbles become too large to contain by its surface pressure, they breakdown bringing about massive shearing force and enormous amount of energy switching from kinetic energy of motion to completely heating of the bubble content. According to Suslick and Doktycz (54), the bubbles have temperature of about 5000 K, pressure of about 1000atm and heating and cooling rate above 1010 K/s. Based on this principle, only liquid and liquid-containing solid materials have cavitations effect. The Ultrasonic assisted extraction can effectively facilitate organic and inorganic compound

leaching from plant matrix. Extraction efficiency is highly affected by moisture content of test sample, milling (processing) degree, particle size, and the nature of solvent used. Likewise, temperature, pressure, recurrence, and time of sonication are the key processing parameters for the activity of ultrasound. The ultrasonic assisted extraction can also be incorporated with other techniques to enhance the efficiency of a conventional system.

The advantages of ultrasonic assisted extraction include the ability to extract at lower temperatures due to the absence of external heat input, low solvent consumption, and reduction in extraction time. Ultrasound energy can likewise cause faster energy transfer, more effective mixing, lower thermal gradients and extraction temperature, specific extraction, and reduced equipment size. Among them, ultrasonic assisted extraction has the remarkable advantages of time and energy savings, higher extraction yield and better quality of essential oils owing to its cavitations, mechanical and thermal effects accelerating the extraction process (55, 56). Foudah *et al.*, (57) reported using the ultrasonic assisted extraction method in extracting cinnamaldehyde from different species of cinnamon. Aryati *et al.*, (58) also extract cinnamaldehyde from *Cinnamomum burmannii* barks.

### Reflux Extraction

Reflux extraction (RE) is a traditional extraction method used to extract bioactive components from plant matrices. Although, thermal degradation at high temperatures can occur over a long period of extraction. Additionally, longer extraction time requires more energy and extraction solvent (59). The procedure for reflux extraction according to the process by Ameer *et al* (60) is done using a water bath-equipped reflux extractor. All extraction experiments were executed utilizing the extraction parameters specified by CCD (Table 2). After extraction, the vessels will be cooled to room temperature and extracts filtered through Whatman filter paper No.41. Afterwards, the obtained extracts will be transferred to falcon tubes and stored at  $4\pm 1$  °C (61). Ahmad *et al.*, (62) and Zhou *et al.*, (20) reported using reflux extraction method to extract cinnamaldehyde as well.

### **Supercritical fluid CO<sub>2</sub> extraction (SFE)**

Supercritical fluids have high solubility of different compounds. Based on this principle, SFE has the advantages of high extraction efficiencies, mild extraction conditions avoiding the decomposition of targeted substances, easy solvent removal without toxic residues, and being environmental-friendly. In SFE, a particular gas is subjected to pressure and temperature past its critical point, above which characteristic gas and liquid phases do not exist. Supercritical fluid has liquid-like thickness and solvation power, and gas-like properties of diffusion, viscosity, and surface strain. These properties make it an excellent solvent to extract flavour and bioactive compounds from plant materials in a short time with higher yields at low temperature. As the pressure decreases, their solvation power increases (63). The most used solvent for SFE is CO<sub>2</sub>, for it possesses the properties of being inert, odourless, inexpensive, non-toxic, non-flammable, and having a comparatively low supercritical pressure (74 bar) and temperature (32 °C). CO<sub>2</sub> is a non-polar molecule, so it is suitable for extracting non-polar compounds. This deficiency can be compensated by using modifiers such as ethanol (64, 65). The SFE is commonly operated in a temperature range of 30-60 °C and a pressure range of 8-40 MPa (52).

It is usually carried out using a supercritical CO<sub>2</sub> extraction apparatus SFE-2 (Applied Separations, America) for cinnamon essential oil extraction. About 20 g ground cinnamon sample will be weighed and then filled into the 50ml vessel. The oven temperature and the exit valve temperature will be set at 60 °C. The one-step static extraction will be performed at 30MPa for 2 hours, with no modifier. The extract will be collected directly without solvents and then sealed for preservation.

The SFE has many advantages over other extraction methods such as the absence of solvent residues and the rich top flavour notes of the extracts. Selective isolation of a single component from plant material can be achieved by using optimal process condition. Main factors influencing the extraction efficiency are temperature, pressure, particle size and moisture content of raw material, extraction time, gas flow rate, and solvent-to-raw material ratio (23). The only disadvantage of

CO<sub>2</sub> is its low polarity, which makes it unsuitable for the extraction of polar substances, which are common in pharmaceuticals and drug samples. This limitation has been successfully overcome using chemical modifier like dichloromethane (CH<sub>2</sub>Cl<sub>2</sub>). Usually, a small amount of modifier can significantly enhance the polarity of carbon dioxide (66). Despite the efficiency in extracting heat-labile bioactive compounds and the use of small quantity of solvent, the use of this method is limited due to the need for high capital and operating cost as a result of the high pressure required for the operation. Many reports have been written on its use to extract cinnamaldehyde from cinnamon species (29, 53, 67-70).

### **Microwave-assisted Hydrodistillation**

In an attempt to take advantage of microwave heating with the conventional hydro distillation, microwave assisted hydro distillation was then developed and used for the extraction of essential oils or extracts from some plants. With this technique, the plant material set in a Clevenger apparatus is heated inside a microwave oven for a brief timeframe to extract the essential oil where heat is produced by microwave energy. The sample reaches its boiling point quickly, resulting in an exceptionally short extraction or distillation time. After extraction, the oil will be dehydrated over anhydrous sodium sulfate to remove excess water, and then the concentrated oil will be weighed and stored in vial at 4 °C. Using the microwave distillation technique, it is possible to achieve distillation with the indigenous water of the fresh plant material (71). This method was also used by Jeyaratnam *et al.* (26). It was done thus Essential oil from *Cinnamomum cassia* (cinnamon) was extracted by using the microwave assisted hydro distillation method. Golmakani and Rezaei (72) also extracted cinnamaldehyde from the cinnamon bark through Microwave assisted hydro distillation. Hearnunyakij and Phutdhawong, (68) extracted cinnamaldehyde using this microwave assisted hydrodistillation and compared it with hydro distillation and super critical fluid extraction method.

### **Validated Pharmacological Properties of Cinnamaldehyde**

Cinnamaldehyde as a predominant component of essential oil extracted from cinnamon have been reported by various scientific investigations to possess therapeutic potentials. Different studies have been carried out to validate the traditional use of cinnamon oil for the management of diseases and infections and also to elucidate the mechanism by which cinnamaldehyde exerts its therapeutic activity. The general mechanism through which cinnamaldehyde exerts its therapeutic potential is through free radical scavenging, prevents the generation of reactive nitrogen species (RNS) or reactive oxygen species (ROS), elevation of cellular antioxidant defense. All these have been implicated in several disease conditions like diabetes, cancer, infectious diseases. Therefore, this section will summarize the various bioactivities of cinnamaldehyde against several infections, disease conditions and boosting general well-being.

#### Antimicrobial activity of cinnamaldehyde

The burden posed by infectious diseases continues to persist despite the different therapeutic agents available for managing infections due to continuous cases of antimicrobial resistance. Bacterial infections account for majority of infections that places both health and economic burden worldwide. *Enterococcus faecium*, *Pseudomonas aeruginosa*, *Klbesiella pneumonia*, *Acinetobacter baumannii*, *Staphylococcus aureus* and *Enterobacter* species are the bacteria of greatest concern and most dangerous human pathogens (73). Cinnamaldehyde has been reported in various studies to possess antimicrobial activities against some food-borne and human pathogens. Due to the continuous multidrug resistance of these pathogens and prevalence of other less threatening and food borne pathogens, there is renewed interest in discovery and development of antibacterial agents from natural sources with low toxicity and affordability. Cinnamaldehyde has also received attention due to the therapeutic potentials folkloric medicine claim to possess. Several literatures have validated the traditional claims and reported the antibacterial activities of cinnamaldehyde against food-borne pathogens and human pathogens.

The antimicrobial activity of cinnamaldehyde in a study carried out by Shen et al., (74)

against gram-positive *Staphylococcus aureus* and gram-negative *Escherichia coli* was observed to be through altering of bacterial cell morphology, disruption of membrane integrity and permeability. In an antivirulence study of cinnamaldehyde against *Staphylococcus aureus* and *Enterococcus faecalis* multidrug resistant strains, it was observed that cinnamaldehyde inhibited the hemolytic activity of the bacteria in human red blood cells. Additionally, the *in vivo* study in *Galleria mellonella* larvae model showed that cinnamaldehyde possess the ability to decrease bacterial virulence and survival rate without causing any phenotypic change (75). Cinnamaldehyde exhibit potent antibacterial activity against various strains of bacteria such as *Streptococcus pyogenes*, *Pseudomonas aeruginosa*, *Staphylococcus epidermidis*, *Staphylococcus pseudintermedius*, *Proteus mirabilis*, *Septococcus mutans* and *Salmonella Typhimurium*, *Aeromonas hydrophila*. Cinnamaldehyde exhibit antimicrobial activity against bacteria by disrupting cell membrane leading to leakage of intracellular components leading eventual death of the cells, inhibiting genes responsible for some proteins and inhibiting the production of biofilms in biofilm producing bacteria. He *et al.*, (76) reported cinnamaldehyde inhibition of biofilm formation by modulating hydrophobicity, aggregation, acid tolerance and acid production. It also inhibited virulent gene expression.

Antifungal potential of cinnamaldehyde in food plants and cash crops has also been explored and reported by numerous studies. Cinnamaldehyde appears to exhibit its antifungal abilities through inhibiting the growth of fungal spores, inhibiting the biosynthesis of aflatoxin in aflatoxin producing fungi *Aspergillus niger*. The aflatoxin biosynthesis is inhibited by altering the normal functions of GPCRs and Oxyipins, biosynthesis of cell wall, as well as redox balance. At a concentration of 0.2  $\mu\text{L/mL}$  Mean inhibitory concentration (MIC) that is, 50% of inhibition was observed in *Alternaria alternata* (77). Some of the fungi that cinnamaldehyde has been reported to possess antifungal activity against include *Aspergillus niger*, *Aspergillus ochraceus*, *Alternaria alternata*, *Candida alicans*, *Malassezia pachydermatis*, *Phytophthora capsici*, *Aspergillus flavus*.

**Table 2.** Summary of mechanism of cinnamaldehyde activities.

Bioactivities	Mechanism	Ref
Antibacterial	Damage bacterial cell morphology Cell membrane integrity and cell permeability damage (Cell lysis) Inhibit biofilm formation	(74, 76,79, 80-82)
Antifungal	Inhibit Salmonella pathogenicity island 1 (SPI-1) Inhibits biosynthesis and degrades mycotoxins Destruction of mitochondria. plasma membrane and loss of cell wall integrity Perturbation of fatty acid, polysaccharide and leucine metabolism	(7, 77, 83-86)
Antiviral	Stimulation of TLR7 pathway Activation and increased expression of IRAK-4	(87)
Antioxidant	Inhibits Aryl hydrocarbon receptor (AHR) activation Activates NRF2/HO1 pathways Up-regulation of Hmox1	(88, 89)
Antidiabetic	Inhibits MAPK signaling pathway Regulates PTP-1B and $\alpha$ -amylase Activates AMPK Upregulates IRS1/PI3K/AKT2 pathway reduced AGE/RAGE interaction	(90-92)
Anti-inflammation	Regulation of Janus kinase/Signal transducer and activator of transcription (Jak/Stat) pathway Blocks NF- $\kappa$ B pathway	(93, 94)
Anticancer	Inhibition of Wnt/ $\beta$ -catenin pathway Inhibit proliferation related protein expression eg. p62 Induction of Beclin-1, ATG5 and LC3B expression	(95, 96)
Anti-neurodegenerative	Inhibits tau aggregation Activation of SIRT1-PGC1 $\alpha$ -PPAR $\gamma$ pathway Reduction of GSK-3 $\beta$ protein level	(97-99)

Cinnamaldehyde has the ability to limit viral infection and replication against enteric viruses. Enteric viruses are viruses that are mainly transmitted through the faecal-oral route, either based on person to person contact or consumption of contaminated food or water. Cinnamaldehyde was observed to have virucidal activity against food-borne pathogens such as human norovirus and hepatitis A virus (78). The mechanisms through which cinnamaldehyde carries out its therapeutic functions are shown in table 2.

### Antioxidant activity of cinnamaldehyde

Oxidative stress caused by the excessive production of reactive oxygen species and reactive nitrogen species (oxidants) in human body accounts for the pathogenesis of a number of diseases such as cancer, diabetes, Alzheimer's disease

(100, 101). The mopping up of these oxidants appears to be a promising measure to reduce the level of oxidative stress. Secondary metabolites known as phytochemicals present in fruits and vegetables, as well as medicinal plants have been reported to scavenge these oxidants (102). Thus, playing an important role in the prevention and management of diseases linked with oxidative stress. Cinnamaldehyde, a potent phenylpropanoid predominant in cinnamon essential oil is majorly responsible for its free radical scavenging and antioxidant activity. The mechanism of action of cinnamaldehyde as an antioxidant agent could be due to up-regulation of antioxidant enzyme Hmox1, reduction in malondialdehyde accumulation and free radical-scavenging. *In vivo* study reported that cinnamaldehyde protected against lipid peroxidation in the brain and kidney of rats (103). All these reports points



cinnamaldehyde as a possible therapeutic agent against oxidative stress related disease conditions.

### Anti-inflammatory activity of cinnamaldehyde

The body has a natural mechanism of defense known as inflammation. This is the process the immune system adopts in recognizing and removing harmful and/or foreign stimuli and initiates the process of healing. Inflammation can be named as acute or chronic relying upon the amount of time it requires to get to a serious state and how long the effects persevere.

Inflammation can be classified as acute or chronic depending on how long it takes to reach a severe state and how long the symptom persists (104). Acute inflammation is an important and constructive response by the immune system against tissue damage however delay in remediation may lead to metabolic related conditions like inflammatory bowel, arthritis and cancer. This occurs as a result of the movement of neutrophils and macrophages into infected tissues from the blood stream (105). Chronic inflammation is a prolonged reaction to biological, chemical or physical stimuli defined by continued deployment of mononuclear leukocytes alongside tissue injury as a result of continual inflammatory response (106). The development and progression of about 25% of cancers have closely been linked to chronic inflammation (107). The shift from synthetic to natural products for managing diseases including diseases associated with inflammation has led to the exploitation of cinnamaldehyde in the bid to develop novel drugs from plants with low toxicity and anti-inflammatory response. Cinnamaldehyde inhibited the IL-8 secretion and expression helicobacter pylori induced gastric inflammation. Additionally, cinnamaldehyde optimally inhibited inflammatory cell infiltration induced by lipopolysaccharide (LPS) (96).

### Antidiabetic activity of cinnamaldehyde

Diabetes is a group of metabolic diseases characterized by high blood glucose level brought about by the alteration in the secretion and/or utilization of insulin. Diabetes is classified into type I and type II diabetes and cinnamaldehyde has been duly documented in literature to possess

antidiabetic properties. Cinnamaldehyde exhibits antihyperglycemic activity in diabetic animal models through increasing the uptake of glucose and enhanced glucogen biosynthesis in the liver, slowing the rate of gastric emptying, pancreatic islets dysfunction restoration and improved diabetic renal and brain disorders. All these effects are achieved by cinnamaldehyde through its action on different signalling pathways, such as Peroxisome proliferated activated receptors (PPARs), AMP-Activated protein kinase (AMPK), GLUT4 and Nrf2 pathways. Nikzamir *et al.*, (108) reported the down regulation of blood glucose through the unregulated expression of GLUT4 gene levels in the skeletal muscle of rodent model. In another study, cinnamaldehyde was observed to have inhibited protein tyrosine phosphatase-1B (PTP-1B), thereby preventing type II diabetes and obesity (90). Cinnamaldehyde boost the antioxidant protection against reactive species under hyperglycemic conditions, which protects the pancreatic beta cells from damages/loss and exhibited hyperglycemic effect (109). In a chemically-induced diabetic rat model, cinnamaldehyde was reported to exhibit hypoglycemic effect against gestational diabetes through modulating PPAR $\gamma$  (110). In a recent study by Zhao *et al.* (111), it was reported that cinnamaldehyde improved glucose metabolism and insulin sensitivity, increased glycogen synthesis in Type 1 diabetes (T1DM). It also regulated the gut microbiota by causing an increase in *Lactobacillus johnsonii* and decrease in *Lactobacillus murinus* in T1DM mice. Glucose-6-phosphate dehydrogenase, an enzyme involved in glucose metabolism was observed to have increased activity because of cinnamaldehyde (112). With the potential of cinnamaldehyde as a possible candidate, there is the need for community based trials and clinical studies to better understand the antidiabetic and hypoglycemic effect of cinnamaldehyde.

### Anti cancer activity of cinnamaldehyde

Cancer is widely referred to as the uncontrolled proliferation of cells that ultimately causes cell aggregation leading the formation of tumours. It is one of the main causes of death yearly. Different factors are linked to the onset of cancers such genetic mutations, smoking, ingestion of pesti-

cides and heavy metals and unhealthy dietary lifestyle. Cellular aggregation from inflammation due to disruption in the proper functioning of signaling pathways eventually results in pathogenesis of cancers. Thus, treatment of cancer depends on the inhibition of cell proliferation and destruction of malicious cells. Selective destruction of tumour cells without affecting normal healthy cells is the target for cancer chemotherapy. Plant sources with anticancer benefits can interfere with possible targets involved with carcinogenesis and tumour cell biology which make them promising tools in cancer research. A number of studies have reported the chemotherapeutic potentials of cinnamaldehyde against various forms of cancers. Various studies have provided scientific proofs of cinnamaldehyde against breast cancer, prostate cancer, lung cancer, colon cancer, hepatocellular carcinoma (113). Cinnamaldehyde exhibits anticancer activity by inducing apoptosis, inhibiting cell viability and proliferation in a dose dependent manner. *In vitro* studies using breast cancer cell lines by Liu *et al* (114) reported inhibition of cell proliferation by cinnamaldehyde with  $IC_{50}$  of 16  $\mu\text{g/ml}$  and 12.23  $\mu\text{g/ml}$  at 24 h and 4 h respectively. *In vitro* and *in vivo* study executed by Chiang *et al.*, (115) reported cinnamaldehyde as the first scientifically demonstrated natural compound to inhibit extracellular and intracellular nicotinamide phosphoribosyl transferase (NAMPT) in reducing visfatin-induced breast cancer progression. In an *in vitro* study, cinnamaldehyde was reported to induce autophagy mediated cell death through the endoplasmic reticulum stress and epigenetic modifications in gastric cancer (96).

### Anti-neurodegenerative activity of cinnamaldehyde

Neurodegenerative diseases are a group of disorders associated with alteration in neuronal normal functions. They are characterized by progressive degeneration and eventual loss of specific neurons. Neurons make up the nervous system which includes the brain and spinal cord and the inability of neurons to reproduce or replace themselves, makes neurodegenerative diseases incurable. They include Alzheimer's disease, Parkinson's disease, Prion disease, Huntington's dis-

ease etc. Cinnamaldehyde has been reported to play important role in reducing neuronal loss and increase the activity of important enzymes in the brain. A study carried out in mouse model of Parkinson's diseases reveals that cinnamaldehyde reduces the dopaminergic neuronal loss in substantia nigra and increases the activity of catalase in the midbrain (116). A major hallmark of Alzheimer's disease is the accumulation of amyloid- $\beta$ . Cinnamaldehyde has been reported to inhibit its formation through its action on the first rate limiting enzyme  $\beta$ -secretase (BACE-1) in amyloid- $\beta$  production. Cinnamaldehyde decrease the level of BACE-1 in 5XFAD transgenic mice (98).

### Applications of Cinnamaldehyde in Health and Medicine

In health and medicine, cinnamaldehyde has proven to influence diabetics in animals and therefore a potential for diabetic intervention in humans (117). It is also applied to traditional medicine to control blood pressure, tumor growth, diabetics, Alzheimer's, and Parkinson's diseases (118). Cinnamaldehyde has the potential to be applied in skin wounds to quicken the healing process (119, 120) and in cancer chemotherapy (113).

### Agriculture

Cinnamaldehyde is applied in the agricultural industry in various ways. It is applied to main the postharvest storage of horticultural crops and maintains the quality of fresh produce (13, 121). It is also used in food preservation as Duan *et al* (122) demonstrated its potential in preserving citrus from green mold. In addition, it is also applied to agriculture for its food preservation and storage quality and to improve the overall quality properties in fruits (123). It is applied in plant breeding to improve the crops architecture for better adaptation to biotic and abiotic stress. Xue *et al* (124) demonstrated this by applying cinnamaldehyde to plants for lateral root formation. Cinnamaldehyde is also applied in agriculture for its potential and use as a pesticide, herbicide, and weedicide (125, 126). It is used in the control of plant diseases and pests in crops in field and during storage of fruits and vegetables (127).

## Food and other Industries

It is applied in the food industry as an effective and promising substitute for chemical synthetic preservative to increase the shelf life of baked food (128). Cinnamaldehyde is also applied as an antifungal agent in food preservation (12). Cinnamaldehyde is applied in the food and beverages, as an odor agent and flavouring agent. Cinnamaldehyde is used in the food industry to reduce the antimicrobial load in packaged food and increase the shelf life of packaged food (129). It is also used as a natural wood preservative (130).

## Conclusion

This review article elucidates the use and

effectiveness of cinnamaldehyde which can be applied to various areas such as medicine, agriculture, food and other industrial purposes. Cinnamaldehyde contains considerable antimicrobial and antifungal properties and can be applied to inhibit the growth of microbes and fungi in food. Cinnamaldehyde has sparked interest in research as its usefulness is seen especially as an anticancer agent. Nevertheless, more studies are required to specify the dosage of cinnamaldehyde for various applications and to also explore other hidden potentials for the benefit of mankind.

## Conflict of Interest

None declared.

## References

1. Almalki DA, Alghamdi SA, Al-Attar AM. Comparative Study on the Influence of Some Medicinal Plants on Diabetes Induced by Streptozotocin in Male Rats. *Biomed Res Int.* 2019 Feb 27;2019:3596287. doi: 10.1155/2019/3596287. PMID: 30937310; PMCID: PMC6415286.
2. Ugboko HU, Nwinyi OC, Oranusi SU, Fatoki TH, Omonhinmin CA. Antimicrobial Importance of Medicinal Plants in Nigeria. *ScientificWorldJournal.* 2020 Sep 22;2020:7059323. doi: 10.1155/2020/7059323. PMID: 33029108; PMCID: PMC7528132.
3. Akinmoladun AC, Ibukun EO, Dan-Ologe IA. Phytochemical constituents and antioxidant properties of extracts from the leaves of *Chromolaena odorata*. *Sci Res Essays.* 2007;2(6):191-194
4. Pathirana H, Wimalasena SHMP, de Silva BCJ, Hossain S, Heo GJ. Antibacterial activity of cinnamon (*Cinnamomum zeylanicum*) essential oil and cinnamaldehyde against fish pathogenic bacteria isolated from cultured olive flounder *Paralichthys olivaceus*. *Indian J Fish.* 2019;66(2):86-92. doi: 10.21077/ijf.2019.66.2.85023-12.
5. Tajkarimi MM, Ibrahim SA, Cliver DO. Antimicrobial herb and spice compounds in food. *Food Control.* 2010;21:1199-1218.
6. Dorri M, Hashemitabar S, Hosseinza deh H. Cinnamon (*Cinnamomum zeylanicum*) as an antidote or a protective agent against natural or chemical toxicities: a review. *Drug Chem Toxicol.* 2018 Jul;41(3):338-351. doi: 10.1080/01480545.2017.1417995. Epub 2018 Jan 10. PMID: 29319361.
7. Wang Y, Feng K, Yang H, Yuan Y, Yue T. Antifungal mechanism of cinnamaldehyde and citral combination against *Penicillium expansum* based on FT-IR fingerprint, plasma membrane, oxidative stress and volatile profile. *RSC Adv.* 2018 Feb 5;8(11):5806-5815. doi: 10.1039/c7ra12191a. PMID: 35539597; PMCID: PMC9078163.
8. Buglak NE, Jiang W, Bahnsen ESM. Cinnamic aldehyde inhibits vascular smooth muscle cell proliferation and neointimal hyperplasia in Zucker Diabetic Fatty rats. *Redox Biol.* 2018 Oct;19:166-178. doi: 10.1016/j.redox.2018.08.013. Epub 2018 Aug 24. PMID: 30172101; PMCID: PMC6122148.
9. Guo W, Wang X, Huang J, Wei C, Wang J, Song L, et al. preparation and antimicrobial effect of cinnamaldehyde-based sustained release fumigant tablet for grain storage. *J Mater Res Technol.* 2020;9(6):14122-14130.
10. Zhang W, Gao J, Cheng C, Zhang M, Liu W, Ma X, et al. Cinnamaldehyde Enhances Antimelanoma Activity through Covalently Binding ENO1 and Exhibits a Promoting Effect with Dacarbazine. *Cancers (Basel).* 2020 Jan 29;12(2):311. doi: 10.3390/cancers12020311. PMID: 32013122; PMCID: PMC7072165.
11. Niu A, Wu H, Ma F, Tan S, Wang G, Qiu W. The antifungal activity of cinnamaldehyde in vapour phase against *Aspergillus niger* isolated from spoiled paddy. *LWT.* 2022;159:113181.
12. Qu S, Yang K, Chen L, Liu M, Geng Q, He X, et al. Cinnamaldehyde, a Promising Natural Preservative Against *Aspergillus flavus*. *Front Microbiol.* 2019 Dec 18;10:2895. doi: 10.3389/

fmicb.2019.02895. PMID: 31921070; PMCID: PMC6930169.

13. Shen Y, Kahramanoğlu I, Chen C, et al. Application of cinnamaldehyde for the postharvest storage of fresh horticultural products. *Horticult Int J*. 2021;5(3):103-105. DOI: 10.15406/hij.2021.05.00212
14. Wang R, Wang R, Yang, B. Extraction of essential oils from five cinnamon leaves and identification of their volatile compound compositions. *Innov. Food Sci Emerg Technol*. 2009;10(2):289-292.
15. Kelly S, Brodie C, Hilkert A. Isotopic-Spectroscopic Technique: Stable Isotope-Ratio Mass Spectrometry (IRMS) In modern techniques for food Authentication, Da-Wen Sun (Ed.). Vol 11. MA, USA.; Academic press, 2018. p.349-413
16. Vangalapati M, Satya S, Prakash S, Avanigadda S, Kiran Sree N, Surya D, et al. A review on pharmacological activities and clinical effects of cinnamon species. *Res J Pharm Biol Chem Sci*. 2012;3(1):653–663.
17. Chen PY, Yu JW, Lu FL, Lin MC, Cheng HF. Differentiating parts of Cinnamomum cassia using LC-qTOF-MS in conjunction with principal component analysis. *Biomed Chromatogr*. 2016 Sep;30(9):1449-57. doi: 10.1002/bmc.3703. Epub 2016 Mar 2. PMID: 26873449.
18. Fan Q, Huang F, Zhou L, He W, Jiang X, Na Z, et al. Development of a strategy for a quick process for separation of volatile compounds in counter-current chromatography: purification of cinnamaldehyde from Cinnamomum cassia by high performance counter-current chromatography. *Prep Biochem Biotechnol*. 2021;51(10):1056-1059. doi: 10.1080/10826068.2021.1894444. Epub 2021 Mar 28. PMID: 33775215.
19. Moreta S, Cahyono E, Affandi N, Fadil F, Kurniawan C. Polymeric and nonpolymeric nanofiber of cinnamaldehyde from cinnamon oil (Cinnamomum Zeylanicum). *J Phys Conf Ser*. 2020;1567 022035.
20. Budiana IGN, Moses KT, and Agus S. Characterization of Cinnamadehyde Compound Isolated from Cinnamon Oil and Its Salmonella Typhy Antibacterial Activity. *J Appl Chem Sci*. 2018;5(2):469-472. <https://dx.doi.org/10.22341/jacs.on.00502p469>
21. Baseri H, Haghighi-Asl, Lotfollahi MN. Effects of Operating Parameters on the Cinnamaldehyde Content of extracted essential Oil Using Various Methods. *Chem Eng Technol*. 2010;33(2):267-274.
22. Yu M, Wang S, Zhu H, Wang H, Yao R, Li F, et al. In-situ reactive heat breaking cell wall by SO<sub>3</sub> hydration: innovative cell-wall breaking technique to enhance extraction of cinnamaldehyde from cinnamon. *Prep Biochem Biotechnol*. 2021;51(9):833-841. doi: 10.1080/10826068.2020.1867867. Epub 2021 Jan 10. PMID: 33427036.
23. Azmir J, Zaidul ISM, Rahman MM, Sharif KM, Mohamed A, Sahena F, et al. Techniques for extraction of bioactive compounds from plant materials: A review. *J Food Engin*. 2013;117:426-436.
24. Aramrueang N, Asavasanti S, Khanunthong A. Leafy Vegetables. In *Integrated Processing Technologies for Food and Agricultural By-products*. Academic Press; 2019. p. 245-272. <https://doi.org/10.1016/B978-0-12-814138-0-00010-1>
25. Kargavkar AM, Patel SB, Pate NB. Extraction of Cinnamon Essential Oil from Cinnamon Bark using Hydro Distillation Method. *Int J Sci Res Dev*. 2017;5(04):663-664
26. Jeyaratnam N, Nour AH, Akindoyo JO. Comparative study between hydrodistillation and microwave-assisted hydrodistillation for extraction of cinnamomum cassia oil. *ARPN J Engin Appl Sci*. 2016;11(4):2647–2652
27. Kasim NN, Ismail SNAS, Masdar ND, Hamid FA, Nawawi WI. Extraction and potential of cinnamon essential oil towards repellency and insecticidal activity. *Int J Sci Res Public*. 2014;4(7):1-6.
28. Vihanova K, Houdkova M, Promgool T, Urbanova K, Kanokmedhakul S, Kokoska L. In-vitro growth-inhibitory effect of essential oils and supercriticalcarbon dioxide extracts from Cinnamomum spp. barks andfruits against food bacterial pathogens in liquid and vaporphase. *J Food Saf*. 2021;e12900.
29. Zhou J, Yuan X, Li L, Zhang T, Wang B. Comparison of different methods for extraction of Cinnamom iramulus: yield, chemical composition and in vitro antiviral activities. *Nat Prod Res*. 2017;31(24):2909-2913, DOI:10.1080/14786419.2017.1299724
30. Alsalm HA, Shawkat MS, Khwaidy M.

- Anitbacterial activity of *Cinnamomum zeylanicum* bark oil and cinnamaldehyde on some locally isolated pathogenic bacteria. *World J Pharm Res.* 2017;6(2):174-185. Doi: 10.20959/wjpr20172-7707
31. Senhaji O, Faid M, Kalalou I. Inactivation of *Escherichia coli* O157:H7 by essential oil from *Cinnamomum zeylanicum*. *Braz J Infect Dis.* 2007 Apr;11(2):234-6. doi: 10.1590/s1413-86702007000200013. PMID: 17625768.
32. Ndlela SC, de Moura JMLN, Olson NK, Johnson LA. Aqueous extraction of oil and protein from soybeans with subcritical water. *J Am Oil Chem Soc.* 2012;89:1145-1153. Doi: 10.1007/s11746-011-1993-7
33. Hassas-Roudsari M, Chang PR, Pegg RB, Tyler RT. Antioxidant capacity of bioactives extracted from canola meal by subcritical water, ethanolic and hot water extraction. *Food Chem.* 2009;114:717-726. doi: 10.1016/j.foodchem.2008.09.097
34. Barbra FJ, Zhu ZZ, Koubaa M. Green alternative methods for the extraction of antioxidant bioactive compounds from winery wastes and by-products: A review. *Trends Food Sci Technol.* 2016;49:96–109. doi: 10.1016/j.tifs.2016.01.006
35. Getachew AT, Cho YG, Chun BS. Effect of pretreatments on isolation of bioactive polysaccharides from spent coffee grounds using subcritical water. *Int J Biol Macromol.* 2018;109:711-719. doi: 10.1016/j.jbiomac.2017.12.120
36. Ko MJ, Kwon HL, Chung MS. Pilot-scale subcritical water extraction of flavonoids from satsuma Mandarin (*Citrus unshiu* Markovich) peel. *Innovative Food Science & Emerging Technologies.* 2016;38:175–181. <https://doi.org/10.1016/j.ifset.2016.10.008>
37. Lee JH, Ko MJ, Chung MS. Subcritical water extraction of bioactive components from red ginseng (*Panax ginseng* C.A. Meyer). *J Supercrit Fluids.* 2018;133:177-183. doi: 10.1016/j.supflu.2017.09.029
38. Nastić N, Švarc-Gajić J, Delerue-Matos C, Barroso MF, Soares C, Moreira MM, et al. Subcritical water extraction as an environmentally-friendly technique to recover bioactive compounds from traditional Serbian medicinal plants. *Ind Crops Prod.* 2018;111:579-589.
39. Pedrasa B, Salema-Oom M, Ša-Nogueirab I, Simõesa P, Paiva A, Barreiros S. Valorization of white wine grape pomace through application of subcritical water: Analysis of extraction, hydrolysis, and biological activity of the extracts obtained. *J Supercrit Fluids.* 2017;128:138–144.
40. Tomšik A, Pavlic B, Vladić J, Cindric M, Jovanov P, Sakac M, et al. Subcritical water extraction of wild garlic (*Allium ursinum* L.) and process optimization by response surface methodology. *J Supercrit Fluids.* 2017;128:79-88. Doi: 10.1016/j.supflu.2017.05.012
41. Yan Z, Luo XP, Cong JL, Zhang HH, Ma HL, Duan YQ. Subcritical water extraction, identification and antiproliferation ability on HepG2 of polyphenols from lotus seed epicarp. *Ind Crops Prod.* 2019;29:472-479. doi: 10.1016/j.indcrop.2018.12.031
42. Jayawardena B, Smith RM. Superheated water extraction of essential oils from *Cinnamomum zeylanicum* (L.). *Phytochem Anal.* 2010 Sep-Oct;21(5):470-2. doi: 10.1002/pca.1221. PMID: 20333616.
43. Eikani MH, Golmohammad F, Sadr ZB, Amoli HS, Mirza M. Optimization of superheated water extraction of essential oils from cinnamon bark using response surface methodology. *J Essent Oil Bear PL.* 2013;16(6):740-748. doi:10.1080/0972060X.2013.813236
44. Guo J, Yang RF, Fan XD, Qiu TQ. Extraction of essential oil from cinnamon by subcritical water. *Chemistry & Industry of Forest Products.* 2014;34(3):92–98.
45. Barrales F M, Rezende CA, Martínez J. Supercritical CO<sub>2</sub> extraction of passion fruit (*Passiflora edulis* sp.) seed oil assisted by ultrasound. *J Supercrit Fluids.* 2015;104:183-192. doi: 10.1016/j.supflu.2015.06.006
46. Hu AJ, Zhang ZH, Zheng J, Wang YM, Chen QX, Liu R, et al. Optimizations and comparison of two supercritical extractions of adlay oil. *Innov Food Sci Emerg Technol.* 2012;13:128–133. <https://doi.org/10.1016/j.fset.2011.10.002>
47. Koubaa M, Mhemdi H, Fages J. Recovery of valuable components and inactivating microorganisms in the agro-food industry with ultrasound-assisted supercritical fluid technology. *J Supercrit Fluids.* 2018;134:71-79.
48. Ma Q, Fan X, Liu X, Qiu T, Jiang J. Ultrasound-enhanced subcritical water extraction of essential oils from *Kaempferia galangal* L. and their comparative antioxidant activities. *Sep Pu-*

*rif Technol.* 2015;150:73-79. doi:10.1016/j.seppur.2015.06.013.

49. Chao Z, Ri-Fun Y, Tai-Qiu Q. Ultrasound-enhanced subcritical water extraction of polysaccharides from *Lycium barbarum* L. *Sep Purif Technol.* 2013;120:141-147. doi: 10.1016/j.seppur.2013.09.044

50. Guo J, Yang R, Gong Y, Hu K, Hu Y, Songa FL. Optimization and evaluation of the ultrasound-enhanced subcritical water extraction of cinnamon bark oil. *LWT.* 2021;141:111673. <https://doi.org/10.1016/j.lwt.2021.111673>

51. De Castro MDL, Garcia-Ayuso LE. Soxhlet extraction of solid materials: An outdated technique with promising innovative future. *Anal Chim Acta.* 1998;369(1-2):1-10. doi: 10.1016/S0003-2670(98)00233-5

52. Ngamwonglumlert L, Devahastin S, Chiewchan N. Natural colorants: Pigment stability and extraction yield enhancement via utilization of appropriate pretreatment and extraction methods. *Crit Rev Food Sci Nutr.* 2017 Oct 13;57(15):3243-3259. doi: 10.1080/10408398.2015.1109498. PMID: 26517806.

53. Masghati S, Ghoreishi SM. (2018). Supercritical CO<sub>2</sub> extraction of cinnamaldehyde and eugenol from cinnamon bark: Optimization of operating conditions via response surface methodology. *J Supercrit Fluids.* 140, 62-71. doi: doi.org/j.supflu.2018.06.002

54. Doktycz SJ, Suslick KS. Interparticle collisions driven by ultrasound. *Science.* 1990;247(4946):1067-1069. Doi: 10.1126/science.2309118

55. Wen C, Zhang J, Zhang H, Dzah CS, Zandile M, Duan Y, Ma H, Luo X. Advances in ultrasound assisted extraction of bioactive compounds from cash crops - A review. *Ultrason Sonochem.* 2018 Nov;48:538-549. doi: 10.1016/j.ultsonch.2018.07.018. Epub 2018 Jul 18. PMID: 30080583.

56. Tekin K, Akalın MK, Şeker MG. Ultrasound bath-assisted extraction of essential oils from clove using central composite design. *Ind Crops Prod.* 2015;77:954-960

57. Foudah AI, Shakeel F, Alqarni MH, Ross SA, Salkini MA, Alam P. Simultaneous Estimation of Cinnamaldehyde and Eugenol in Essential Oils and Traditional and Ultrasound-Assisted Extracts of Different Species of Cinnamon Using

a Sustainable/Green HPTLC Technique. *Molecules.* 2021 Apr 3;26(7):2054. doi: 10.3390/molecules26072054. PMID: 33916710; PMCID: PMC8038348.

58. Aryati WD, Nadhira A, Febianli D, Fransisca F, Mun'im A. Natural deep eutectic solvents ultrasound-assisted extraction (NADES-UAE) of trans-cinnamaldehyde and coumarin from cinnamon bark [*Cinnamomum burmannii* (Nees & T. Nees) Blume]. *J Res Pharm.* 2020;24(3):389-398. doi: 10.35333/jrp.2020.162

59. Ameer K, Shahbaz HM, Kwon JH. Green extraction methods for polyphenols from plant matrices and their byproducts: A review. *Compr Rev Food Sci. F.* 2017;16(2): 295-315. doi: 10.1011/1541-4337.12253.

60. Ameer K, Bae SW, Jo Y, Chung N, Gao Y, Kwon JH. Optimization and modeling for heat reflux extraction of total yield, stevioside and rebaudioside-A from *Stevia rebaudiana* (Bertoni) leaves. *Sep Sci Technol.* 2017;52(7):1193-1205 <https://doi.org/10.1080/01496395.2017.1285313>

61. Lee HG, Jo Y, Ameer K, Kwon JH. Optimization of green extraction methods for cinnamic acid and cinnamaldehyde from Cinnamon (*Cinnamomum cassia*) by response surface methodology. *Food Sci Biotechnol.* 2018 Jul 27;27(6):1607-1617. doi: 10.1007/s10068-018-0441-y. PMID: 30483424; PMCID: PMC6233399.

62. Ahmad I, Arifianti AE, Sakti AS, Saputri FC, Mun'im A. Simultaneous Natural Deep Eutectic Solvent-Based Ultrasonic-Assisted Extraction of Bioactive Compounds of Cinnamon Bark and Sappan Wood as a Dipeptidyl Peptidase IV Inhibitor. *Molecules.* 2020 Aug 23;25(17):3832. doi: 10.3390/molecules25173832. PMID: 32842548; PMCID: PMC7504365.

63. Sowbhagya HB, Chitra VN. Enzyme-assisted extraction of flavorings and colorants from plant materials. *Crit Rev Food Sci Nutr.* 2010 Feb;50(2):146-61. doi: 10.1080/10408390802248775. PMID: 20112157.

64. Hamburger M, Baumann D, Adler S. Supercritical carbon dioxide extraction of selected medicinal plants--effects of high pressure and added ethanol on yield of extracted substances. *Phytochem Anal.* 2004 Jan-Feb;15(1):46-54. doi: 10.1002/pca.743. PMID: 14979527.

65. Akanda MJ, Sarker MZ, Ferdosh S, Manap MY, Ab Rahman NN, Ab Kadir MO. Ap-

plications of supercritical fluid extraction (SFE) of palm oil and oil from natural sources. *Molecules*. 2012 Feb 10;17(2):1764-94. doi: 10.3390/molecules17021764. PMID: 22328076; PMCID: PMC6268233.

66. Hawthorne SB, Miller DJ. Direct comparison of soxhlet and low-and high temperature supercritical CO<sub>2</sub> extraction efficiencies of organics from environmental solids. *Anal Chem*. 1994;66:4005-4012.

67. Wang L, Jin J, Liu X, Wang Y, Liu Y, Zhao Y, Xing F. Effect of Cinnamaldehyde on Morphological Alterations of *Aspergillus ochraceus* and Expression of Key Genes Involved in Ochratoxin A Biosynthesis. *Toxins (Basel)*. 2018 Aug 22;10(9):340. doi: 10.3390/toxins10090340. PMID: 30135391; PMCID: PMC6162615.

68. Hearunyakij M, Phutdhawong W. (2022). The Process Optimization Of Microwave Assisted Extraction On Cinnamomum Verum J. Presl. Extraction Yield And Comparative Of Chemical Compositions To Supercritical Fluid Extraction And Hydrodistillation. *Nat Volatiles & Essent. Oils*. 2022;9(1):1343-1358

69. Marongiu B, Piras A, Porcedda S, Tuveri E, Sanjust E, Meli M, Sollai F, Zucca P, Rescigno A. Supercritical CO<sub>2</sub> extract of *Cinnamomum zeylanicum*: chemical characterization and antityrosinase activity. *J Agric Food Chem*. 2007 Nov 28;55(24):10022-7. doi: 10.1021/jf071938f. Epub 2007 Oct 30. PMID: 17966976.

70. Jahangir MA, Shehzad A, Butt MS, Shahid B. Influence of supercritical fluid extract of *Cinnamomum zeylanicum* bark on physical, bioactive and sensory properties of innovative cinnamaldehyde-enriched chocolates. *Czech J Food Sci*. 2018;36: 28-36. doi:10.17221/237/2016-CJFS

71. Kürkcüoğlu M, Baser KHC. Headspace volatiles of three Turkish plants. *J Essent Oil Res*. 2010;22(5):389-392.

72. Golmakani MT, Rezaei K. Comparison of microwave-assisted hydrodistillation with the traditional hydrodistillation method in the extraction of essential oils from *Thymus vulgaris* L. *Food Chem*. 2008 Aug 15;109(4):925-30. doi: 10.1016/j.foodchem.2007.12.084. Epub 2008 Jan 26. PMID: 26050009.

73. Cano A, Etcheto M, Espina M, López-Machado A, Cajal Y, Rabanal F, Sánchez-López E, Camins A, García ML, Souto EB. State-of-the-

art polymeric nanoparticles as promising therapeutic tools against human bacterial infections. *J Nanobiotechnology*. 2020 Oct 31;18(1):156. doi: 10.1186/s12951-020-00714-2. PMID: 33129333; PMCID: PMC7603693.

74. Shen S, Zhang T, Yuan Y, Lin S, Xu J, Ye H. Effects of cinnamaldehyde on *Escherichia coli* and *Staphylococcus aureus* membrane. *Food Control*. 2015;47:196-202. doi:10.1016/j.foodcont.2014.07.003.

75. Ferro TA, Araújo JM, Dos Santos Pinto BL, Dos Santos JS, Souza EB, da Silva BL, et al. Cinnamaldehyde Inhibits *Staphylococcus aureus* Virulence Factors and Protects against Infection in a *Galleria mellonella* Model. *Front Microbiol*. 2016 Dec 21;7:2052. doi: 10.3389/fmicb.2016.02052. Erratum in: *Front Microbiol*. 2021 Jan 15;11:628074. PMID: 28066373; PMCID: PMC5174152.

76. He Z, Huang Z, Jiang W, Zhou W. Antimicrobial Activity of Cinnamaldehyde on *Streptococcus mutans* Biofilms. *Front Microbiol*. 2019 Sep 25;10:2241. doi: 10.3389/fmicb.2019.02241. PMID: 31608045; PMCID: PMC6773874.

77. Xu L, Tao N, Yang W, Jing G. Cinnamaldehyde damaged the cell membrane of *Alternaria alternata* and induced the degradation of mycotoxins in vivo. *Ind Crops Prod*. 2018;112:427-433. doi:10.1016/j.indcrop.2017.12.038

78. Fabra MJ, Castro-Mayorga JL, Randazzo W, Lagarón JM, López-Rubio A, Aznar R, et al. Efficacy of Cinnamaldehyde Against Enteric Viruses and Its Activity After Incorporation Into Biodegradable Multilayer Systems of Interest in Food Packaging. *Food Environ Virol*. 2016 Jun;8(2):125-32. doi: 10.1007/s12560-016-9235-7. Epub 2016 Mar 23. PMID: 27008344.

79. Firmino DF, Cavalcante TTA, Gomes GA, Firmino NCS, Rosa LD, de Carvalho MG, Catunda FEA Jr. Antibacterial and Antibiofilm Activities of *Cinnamomum* Sp. Essential Oil and Cinnamaldehyde: Antimicrobial Activities. *ScientificWorldJournal*. 2018 Jun 6;2018:7405736. doi: 10.1155/2018/7405736. PMID: 29977171; PMCID: PMC6011056.

80. Albano M, Crulhas BP, Alves FCB, Pereira AFM, Andrade BFMT, Barbosa LN, et al. Antibacterial and anti-biofilm activities of cinnamaldehyde against *S. epidermidis*. *Microb Pathog*. 2019 Jan;126:231-238. doi: 10.1016/j.

micpath.2018.11.009. Epub 2018 Nov 12. PMID: 30439400.

81. Liu Y, Zhang Y, Zhou Y, Wang T, Deng X, Chu X, et al. Cinnamaldehyde inhibits type three secretion system in *Salmonella enterica* serovar Typhimurium by affecting the expression of key effector proteins. *Vet Microbiol.* 2019 Dec;239:108463. doi: 10.1016/j.vetmic.2019.108463. Epub 2019 Oct 21. PMID: 31767076.

82. Yin L, Chen J, Wang K, Geng Y, Lai W, Huang X, et al. Study the antibacterial mechanism of cinnamaldehyde against drug-resistant *Aeromonas hydrophila* in vitro. *Microb Pathog.* 2020 Aug;145:104208. doi: 10.1016/j.micpath.2020.104208. Epub 2020 Apr 20. PMID: 32325237.

83. Sim JXF, Khazandi M, Pi H, Venter H, Trott DJ, Deo P. Antimicrobial effects of cinnamon essential oil and cinnamaldehyde combined with EDTA against canine otitis externa pathogens. *J Appl Microbiol.* 2019 Jul;127(1):99-108. doi: 10.1111/jam.14298. Epub 2019 May 21. PMID: 31050849.

84. Sun Q, Shang B, Wang L, Lu Z, Liu Y. Cinnamaldehyde inhibits fungal growth and aflatoxin B1 biosynthesis by modulating the oxidative stress response of *Aspergillus flavus*. *Appl Microbiol Biotechnol.* 2016 Feb;100(3):1355-1364. doi: 10.1007/s00253-015-7159-z. Epub 2015 Nov 19. PMID: 26585445.

85. Wang P, Ma L, Jin J, Zheng M, Pan L, Zhao Y, et al. The anti-aflatoxicogenic mechanism of cinnamaldehyde in *Aspergillus flavus*. *Sci Rep.* 2019 Jul 19;9(1):10499. doi: 10.1038/s41598-019-47003-z. PMID: 31324857; PMCID: PMC6642104.

86. Wang Y, Wang M, Li M, Zhao T, Zhou L. Cinnamaldehyde inhibits the growth of *Phytophthora capsici* through disturbing metabolic homeostasis. *PeerJ.* 2021 Apr 30;9:e11339. doi: 10.7717/peerj.11339. PMID: 33987017; PMCID: PMC8092109.

87. Liu R, He T, Zeng N, Chen T, Gou L, Liu J. Mechanism of anti-influenza virus of cinnamaldehyde and volatile oils from *Ramulus cinnamomi*. *Med plant.* 2015;6(1-2):4-8

88. Uchi H, Yasumatsu M, Morino-Koga S, Mitoma C, Furue M. Inhibition of aryl hydrocarbon receptor signaling and induction of NRF2-mediated antioxidant activity by cinnamaldehyde in human keratinocytes. *J Dermatol Sci.* 2017 Jan;85(1):36-43. doi: 10.1016/j.jdermsci.2016.10.003. Epub 2016 Oct 4. PMID: 27720465.

89. Tanaka Y, Uchi H, Furue M. (2019). Antioxidant cinnamaldehyde attenuates UVB-induced photoaging. *J Dermatol Sci.* 2019;96(3):151-158. doi:10.1016/j.jdermsci.2019.11.001

90. Saifudin A, Kadota S, Tezuka Y. Protein tyrosine phosphatase 1B inhibitory activity of Indonesian herbal medicines and constituents of *Cinnamomum burmannii* and *Zingiber aromaticum*. *J Nat Med.* 2013 Apr;67(2):264-70. doi: 10.1007/s11418-012-0674-7. Epub 2012 May 30. PMID: 22645080.

91. Gao J, Zhang M, Niu R, Gu X, Hao E, Hou X, et al. The combination of cinnamaldehyde and kaempferol ameliorates glucose and lipid metabolism disorders by enhancing lipid metabolism via AMPK activation. *J Funct Foods.* 2021;83:104556. doi: 10.1016/j.jff.2021.104556

92. Abdelmageed ME, Shehatou GS, Abdelsalam RA, Suddek GM, Salem HA. Cinnamaldehyde ameliorates STZ-induced rat diabetes through modulation of IRS1/PI3K/AKT2 pathway and AGEs/RAGE interaction. *Naunyn Schmiedebergs Arch Pharmacol.* 2019 Feb;392(2):243-258. doi: 10.1007/s00210-018-1583-4. Epub 2018 Nov 20. PMID: 30460386.

93. Cheng WX, Zhong S, Meng XB, Zheng NY, Zhang P, Wang Y, Qin L, Wang XL. Cinnamaldehyde Inhibits Inflammation of Human Synovial Cells Through Regulation of Jak/Stat Pathway and Ameliorates Collagen-Induced Arthritis in Rats. *J Pharmacol Exp Ther.* 2020 May;373(2):302-310. doi: 10.1124/jpet.119.262907. Epub 2020 Feb 6. PMID: 32029577.

94. Muhammad JS, Zaidi SF, Shaharyar S, Refaat A, Usmanhany K, Saiki I, Sugiyama T. Anti-inflammatory effect of cinnamaldehyde in *Helicobacter pylori* induced gastric inflammation. *Biol Pharm Bull.* 2015;38(1):109-15. doi: 10.1248/bpb.b14-00609. PMID: 25744466.

95. Wu C, Zhuang Y, Jiang S, Tian F, Teng Y, Chen X, et al. Cinnamaldehyde induces apoptosis and reverses epithelial-mesenchymal transition through inhibition of Wnt/ $\beta$ -catenin pathway in non-small cell lung cancer. *Int J Biochem Cell Biol.* 2017 Mar;84:58-74. doi: 10.1016/j.biocel.2017.01.005. Epub 2017 Jan 16. PMID:



28093328.

96. Kim TW. Cinnamaldehyde induces autophagy-mediated cell death through ER stress and epigenetic modification in gastric cancer cells. *Acta Pharmacol Sin.* 2022 Mar;43(3):712-723. doi: 10.1038/s41401-021-00672-x. Epub 2021 May 12. PMID: 33980998; PMCID: PMC8888591.

97. Peterson DW, George RC, Scaramozzino F, LaPointe NE, Anderson RA, Graves DJ, et al. Cinnamon extract inhibits tau aggregation associated with Alzheimer's disease in vitro. *J Alzheimers Dis.* 2009;17(3):585-97. doi: 10.3233/JAD-2009-1083. PMID: 19433898.

98. Do J, Kim N, Jeon SH, Gee MS, Ju YJ, Kim JH, Oh MS, Lee JK. Trans-Cinnamaldehyde Alleviates Amyloid-Beta Pathogenesis via the SIRT1-PGC1 $\alpha$ -PPAR $\gamma$  Pathway in 5XFAD Transgenic Mice. *Int J Mol Sci.* 2020 Jun 24;21(12):4492. doi: 10.3390/ijms21124492. PMID: 32599846; PMCID: PMC7352815.

99. Emamghoreishi M, Farrokhi MR, Amiri A, Keshavarz M. The neuroprotective mechanism of cinnamaldehyde against amyloid- $\beta$  in neuronal SHSY5Y cell line: The role of N-methyl-D-aspartate, ryanodine, and adenosine receptors and glycogen synthase kinase-3 $\beta$ . *Avicenna J Phytomed.* 2019 May-Jun;9(3):271-280. PMID: 31143694; PMCID: PMC6526042.

100. Zhang YJ, Gan RY, Li S, Zhou Y, Li AN, Xu DP, et al. Antioxidant Phytochemicals for the Prevention and Treatment of Chronic Diseases. *Molecules.* 2015 Nov 27;20(12):21138-56. doi: 10.3390/molecules201219753. PMID: 26633317; PMCID: PMC6331972.

101. Ahmad KA, Yuan Yuan D, Nawaz W, Ze H, Zhuo CX, Talal B, et al. Antioxidant therapy for management of oxidative stress induced hypertension. *Free Radic Res.* 2017 Apr;51(4):428-438. doi: 10.1080/10715762.2017.1322205. Epub 2017 May 9. PMID: 28427291.

102. Chopra B, Dhingra AK, Dhar KL. *Psoralea corylifolia* L. (Buguchi) - folklore to modern evidence: review. *Fitoterapia.* 2013 Oct;90:44-56. doi: 10.1016/j.fitote.2013.06.016. Epub 2013 Jul 4. PMID: 23831482.

103. Sharma UK, Sharma AK, Gupta A, Kumar R, Pandey A, Pandey AK. Pharmacological activities of cinnamaldehyde and eugenol: antioxidant, cytotoxic and anti-leishmanial studies. *Cell Mol Biol (Noisy-le-grand).* 2017 Jul 31;63(6):73-

78. doi: 10.14715/cmb/2017.63.6.15. PMID: 28968213.

104. Germolec DR, Shipkowski KA, Frawley RP, Evans E. (2018). Markers of Inflammation. In: DeWitt, J., Rockwell, C., Bowman, C. (eds) Immunotoxicity Testing. Methods in Molecular Biology. 2018; Vol. 1803. Humana Press, New York, NY. doi:10.1007/978-1-4939-8549-4\_5

105. Ferrero-Miliani L, Nielsen OH, Andersen PS, Girardin SE. Chronic inflammation: importance of NOD2 and NALP3 in interleukin-1beta generation. *Clin Exp Immunol.* 2007 Feb;147(2):227-35. doi: 10.1111/j.1365-2249.2006.03261.x. PMID: 17223962; PMCID: PMC1810472.

106. Fleit HB. Chronic inflammation. Pathobiology of Human Disease: A dynamic Encyclopedia of disease mechanisms. Stony Brook University School of medicine, Stony Brook, NY, USA. 2014; p. 300-314. doi:10.16/B978-0-12-386456-7.018-8-6

107. Hussain SP, Harris CC. Inflammation and cancer: an ancient link with novel potentials. *Int J Cancer.* 2007 Dec 1;121(11):2373-80. doi: 10.1002/ijc.23173. PMID: 17893866.

108. Nikzamir A, Palangi A, Kheirollaha A, Tabar H, Malakaskar A, Shahbazian H, et al. Expression of Glucose Transporter 4 (GLUT4) is Increased by Cinnamaldehyde in C2C12 Mouse Muscle Cells. *Iran Red Crescent Med J.* 2014 Feb;16(2):e13426. doi: 10.5812/ircmj.13426. Epub 2014 Feb 4. PMID: 24719730; PMCID: PMC3965863.

109. Subash-Babu P, Alshatwi AA, Ignacimuthu S. Beneficial Antioxidative and Antiperoxidative Effect of Cinnamaldehyde Protect Streptozotocin-Induced Pancreatic  $\beta$ -Cells Damage in Wistar Rats. *Biomol Ther (Seoul).* 2014 Jan;22(1):47-54. doi: 10.4062/biomolther.2013.100. PMID: 24596621; PMCID: PMC3936432.

110. Hosni AA, Abdel-Moneim AA, Abdel-Reheim ES, Mohamed SM, Helmy H. Cinnamaldehyde potentially attenuates gestational hyperglycemia in rats through modulation of PPAR $\gamma$ , proinflammatory cytokines and oxidative stress. *Biomed Pharmacother.* 2017 Apr;88:52-60. doi: 10.1016/j.biopha.2017.01.054. Epub 2017 Jan 13. PMID: 28092845.

111. Zhao H, Wu H, Duan M, Liu R, Zhu Q, Zhang K, Wang L. Cinnamaldehyde Improves Metabolic Functions in Streptozotocin-Induced

- Diabetic Mice by Regulating Gut Microbiota. *Drug Des Devel Ther.* 2021 Jun 1;15:2339-2355. doi: 10.2147/DDDT.S288011. PMID: 34103897; PMCID: PMC8179756.
112. Çelik Z, Mert H, Comba B, Mert N. Effects of cinnamaldehyde on glucose-6-phosphate dehydrogenase activity, some biochemical and hematological parameters in diabetic rats. *Biomarkers.* 2022;27(3):270-277. doi: 10.1080/1354750X.2022.2032351
113. Hong SH, Ismail IA, Kang SM, Han DC, Kwon BM. Cinnamaldehydes in Cancer Chemotherapy. *Phytother Res.* 2016;30(5):754-767. doi:10.1002/ptr.5592
114. Liu Y, An T, Wan D, Yu B, Fan Y, Pei X. Targets and Mechanism Used by Cinnamaldehyde, the Main Active Ingredient in Cinnamon, in the Treatment of Breast Cancer. *Front Pharmacol.* 2020 Dec 9;11:582719. doi: 10.3389/fphar.2020.582719. PMID: 33536908; PMCID: PMC7848847.
115. Chiang YF, Chen HY, Huang KC, Lin PH, Hsia SM. Dietary Antioxidant Trans-Cinnamaldehyde Reduced Visfatin-Induced Breast Cancer Progression: In Vivo and In Vitro Study. *Antioxidants (Basel).* 2019 Dec 6;8(12):625. doi: 10.3390/antiox8120625. PMID: 31817697; PMCID: PMC6943554.
116. Mehraein F, Zamani M, Negahdar F, Shojaee A. Cinnamaldehyde attenuates dopaminergic neuronal loss in substantia nigra and induces midbrain catalase activity in a mouse model of Parkinson's disease. *J Basic Clin Pathophysiol.* 2018; 6(1):9-16.
117. Zhu R, Liu H, Liu C, Wang L, Ma R, Chen B, et al. Cinnamaldehyde in diabetes: A review of pharmacology, pharmacokinetics and safety. *Pharmacol Res.* 2017 Aug;122:78-89. doi: 10.1016/j.phrs.2017.05.019. Epub 2017 May 27. PMID: 28559210.
118. Ribeiro-Santos R, Andrade M, Madella D, Martinazzo AP, Moura LAG, de Melo NR, et al. Revisiting an ancient spice with medicinal purposes: Cinnamon. *Trends Food Sci Technol.* 2017;62:154-169. doi:10.1016/j.tifs.2017.02.011
119. Ferro TAF, Souza EB, Suarez MAM, Rodrigues JFS, Pereira DMS, Mendes SJF, et al. Topical Application of Cinnamaldehyde Promotes Faster Healing of Skin Wounds Infected with *Pseudomonas aeruginosa*. *Molecules.* 2019 Apr 25;24(8):1627. doi: 10.3390/molecules24081627. PMID: 31027179; PMCID: PMC6515316.
120. Kucinska-lipka J, Gubanska I, Lewandowska A, Terebieniec A, Przybytek A, Cieśliński. Antibacterial polyurethanes modified with cinnamaldehyde, as potential materials for fabrication of wound dressings. *Polym Bull.* 2019;75:2725-2742. doi: 10.1007/s00289-018-2512-x
121. Shen Y, Kahramanoğlu I, Chen C, Chen J, Okatan V, Wan C. Application of cinnamaldehyde for postharvest storage of fresh horticultural products. *Horticult Int J.* 2021;5(3):103-105. Doi:10.15406/hij.2021.05.00212
122. Duan X, OuYang Q, Tao N. Effect of applying cinnamaldehyde incorporated in wax on green mould decay in citrus fruits. *J Sci Food Agric.* 2018;98(2):527-533. doi: 10.1002/jsfa.8490.
123. Xu Y, Yin Y, Li T, Zhao H, Li X, Li J, et al. Effects of lysozyme combined with cinnamaldehyde on storage quality of olive flounder (*Paralichthys olivaceus*) fillet. *J Food Sci.* 2020;85(4):1037-1044. doi: 10.1111/1750-3841.14980
124. Xue YF, Zhang M, Qi ZQ, Li YQ, Shi Z, Chen J. Cinnamaldehyde promotes root branching by regulating endogenous hydrogen sulfide. *J Sci Food Agric.* 2016 Feb;96(3):909-14. doi: 10.1002/jsfa.7164. Epub 2015 Mar 27. PMID: 25752512.
125. Lu L, Shu C, Chen L, Yang Y, Ma S, Zhu K, et al. Insecticidal activity and mechanism of cinnamaldehyde in *C. elegans*. *Fitoterapia.* 2020 Oct;146:104687. doi: 10.1016/j.fitote.2020.104687. Epub 2020 Jul 15. PMID: 32681860.
126. Muñoz M, Torres-Pagán, Peiro R, Guisjarro R, Sánchez-Moreiras AM, Verdeguer M. phytotoxic effects of three natural compounds: Pelargononic acid, carvacrol and cinnamic aldehyde, against problematic weeds in Mediterranean crops. *Agronomy.* 2020;10(6):791. doi:10.3390/agronomy10060791
127. Kowalska J, Tyburski J, Matysiak K, Jakubowska M, Łukaszyk J, Krzysińska J. Cinnamon as a Useful Preventive Substance for the Care of Human and Plant Health. *Molecules.* 2021 Aug 31;26(17):5299. doi: 10.3390/molecules26175299. PMID: 34500731; PMCID: PMC8433798.
128. Sun Q, Li J, Sun Y, Chen Q, Zhang L, Le T. The antifungal effects of cinnamaldehyde against

*Aspergillus niger* and its application in bread preservation. *Food Chem.* 2020;317:126405. doi: 10.1016/j.foodchem.2020.126405

129. Olszewska MA, Gędas A, Simões M. The Effects of Eugenol, Trans-Cinnamaldehyde, Citronellol, and Terpineol on *Escherichia coli* Biofilm Control as Assessed by Culture-Dependent

and -Independent Methods. *Molecules.* 2020 Jun 6;25(11):2641. doi: 10.3390/molecules25112641. 130. Fang S, Feng X, Lei Y, Chen Z, Yan L.

Improvement of wood decay resistance with cinnamaldehyde chitosan emulsion. *Ind Crops Prod.* 2021;160:113118. doi: 10.1016/j.indcrop.2020.113118

